

Pion re-scattering in π^0 production near threshold

V. Malafaia^{a *} and M. T. Peña^{a†}

^aCFIF and Department of Physics, Instituto Superior Técnico,
1049-001 Lisboa, Portugal

We compare the most used approximations to calculate the contribution from pion re-scattering mechanisms to pion production from proton-proton scattering, near threshold. Our calculation employs a pseudo-vector coupling for the πNN vertex and realistic amplitudes for the πN re-scattering and the NN transitions.

1. Introduction

Due to isospin suppression of the otherwise dominant low-energy isovector Weinberg-Tomozawa πN term in the re-scattering diagram, and the negative interference between the remaining impulse and isoscalar pion re-scattering terms[1], the empirical data for $pp \rightarrow pp\pi^0$ near threshold is only explained by phenomenological non-pionic, short-range, two-body mechanisms[2], or alternatively by an off-shell extrapolation of the πN amplitude[3]. The high quality data for $pp \rightarrow pp\pi^0$ becomes therefore specially interesting by allowing to establish the importance of the short-range mechanisms, which are difficult to constrain: they are subsumed in the low energy constants appearing also in the two- and three-nucleon potentials, but so far these constants have not yet been precisely determined[4].

Nevertheless, any conclusion to be read off from the π^0 production data is only meaningful and quantitative provided that the calculation of the re-scattering diagram[5] is under absolute control. This is not the case in calculations performed till now, where approximations at the level of the energy integration of the corresponding Feynman diagram were considered. Namely, non-relativistic reductions of the Feynman π -exchange diagram were done, through choices for the energy of the exchanged pion, both for its propagator and for the $\pi N - \pi N$ amplitude at the re-scattering vertex.

This work investigates the validity of the mostly used approximations. It generalizes the work of Refs.[5] which considers a toy model for scalar particles and interactions. Our calculation employs a physical model for nucleons and pions, with a pseudo-vector coupling for the πNN vertex; the χ Pt $\pi N - \pi N$ re-scattering amplitude[1] and the Bonn B potential for the nucleon-nucleon interaction in the final state.

*Work supported by FCT under the grant SFRD/BD/4876/2001.

†Work supported by FCT under the grant CERN/FIS/43709/2001.

2. From the Feynman diagram to DWBA

Starting with the Feynman diagram, the energy integration in the energy Q'_0 of the exchanged pion was done after a partial fraction decomposition, which isolates the poles of the pion propagator. This way we separated the DWBA amplitude from the stretched boxes (where two exchanged particles may be in flight simultaneously) contribution. Then,

$$\mathcal{M}_{DWBA} = \frac{1}{2} \int \frac{d^3 q'}{(2\pi)^3} \left[V(\omega_\pi) G_\pi \frac{(\omega_1 - \omega_2) + E_\pi}{\omega_\pi} \right] \frac{1}{(E_1 + E_2 - E_\pi - \omega_1 - \omega_2)} T_{NN}^{FSI} \quad (1)$$

is the exact expression for the DWBA amplitude, where T stands for the transition-matrix of the final-state interaction and $V(\omega_\pi)$ is the product of the $\pi N - \pi N$ amplitude with the πNN vertex. The pion propagator turned to be exactly given by

$$G_\pi = \frac{1}{\left[\frac{\omega_1 - \omega_2}{2} + \frac{E_\pi}{2} \right]^2 - \left[\left(E_{tot} - E - \frac{E_\pi}{2} \right) - \frac{\omega_1 + \omega_2}{2} - \omega_\pi \right]^2}, \quad (2)$$

Thus, Eq.(1) is taken as the reference result, in order to investigate the effect of the most common approximations hitherto used for the pion propagator in a DWBA final-state calculation. These approximations correspond to the following three different choices:

$$\begin{aligned} G_\pi^{fk} &\equiv \frac{1}{\left(\frac{m_\pi}{2} \right)^2 - \omega_\pi^2} && \text{fixed kinematics} \\ G_\pi^{on} &\equiv \frac{1}{(E - \omega_2)^2 - \omega_\pi^2} && \text{on-shell} \\ G_\pi^{st} &\equiv -\frac{1}{\omega_\pi^2} && \text{static approximation} \end{aligned} \quad (3)$$

The “fixed kinematics” approximation “freezes” the energy for the exchanged pion as the difference between the two on-mass-shell energies of the nucleon, before and after re-scattering, at $m_\pi/2$, its value at threshold; the “on-shell approximation” considers for the energy of the pion the difference between the same on-mass-shell energies of the nucleon, but it varies with the energy available for the production process, and is not frozen at the threshold point. Naturally, the two approximations coincide at threshold and the deviation between them increases with energy. The “static approximation” makes the pion exchange instantaneous and therefore neglects energy transfer. It is part of the usual non-relativistic approximation for exchange Feynman diagrams.

3. Results and Conclusions

For the toy model case of scalar particles and scalar interactions, we verified the dominance of the DWBA amplitude over the stretched boxes amplitude, as obtained in Ref.[5]. Furthermore, our calculations show that the DWBA keeps being dominant, as seen on the left panel of Fig.1, even in a more pronounced way for the physical case considered here. We found also that within the scalar toy model all the approximations for the pion energy taken at the $\pi N - \pi N$ re-scattering amplitude overestimate the cross section, as found in Refs.[5]. However, on the right panel of Fig.1, where we describe the nucleon-nucleon interaction by the Bonn B potential T-matrix, only the on-shell G^{on} approximation to the exact pion propagator gives a larger cross section (on the right panel of Fig.1 this curve practically coincides with the reference result). The choice for the energy in the




Figure 1. Left: The stretched boxes compared to DWBA (FSI) for the cross section as a function of the mass of the exchanged particle. Right: Effect of the choices for the energy of the exchanged pion. The nucleon-nucleon interaction is the Bonn B potential.

pion propagator is not very decisive, justifying the usual low-energy static or instantaneous approximation for exchange Feynman diagrams. However, the choice for the energy of the exchanged pion for the re-scattering vertex, as prescribed from the 4-dimensional Feynman diagram, is crucial for the cross section strength. As shown on the right panel of Fig. 1, the fixed kinematics and the on-shell prescriptions for the re-scattering vertex are not suitable for combination with an energy independent NN potential.

Both for the final- and initial-state interaction, our results show that the DWBA formalism is quite adequate at threshold since this part of the full amplitude is clearly dominant over the stretched boxes. Also, the effect of the usual choices for the πN re-scattering energy, which is not fixed by a non-relativistic formalism, can be significant.

In order to extract quantitatively from the data the unknown strength of the short-range mechanisms for the process $pp \rightarrow pp\pi^0$, one has to use for the pion exchange diagram the reference amplitudes. The latter were obtained as non-relativistic reductions of the corresponding Feynman diagrams for the final- and initial-state interactions, respectively.

REFERENCES

1. Y. Park et al., Phys. Rev. **C70**, 1519 (1996); T. D. Cohen et al., Phys. Rev. **C53**, 2661 (1996); T. Sato et al., Phys. Rev. **C56**, 1246 (1997).
2. T.-S. Lee and D. O. Riska, Phys. Rev. Lett. **70**, 2237 (1993); U. van Kolck et al., Phys. Lett. **B388**, 679 (1996); C. Hanhart et al., Phys. Lett **B444** 25 (1998); M. T. Peña et al., Phys. Rev. **C60**, 045201 (1999).
3. E. Hernández and E. Oset, Phys. Lett. **B350** 158 (1995).
4. E. Epelbaum et al., Eur. Phys. J. **A15**, 543 (2002).
5. C. Hanhart et al., Phys. Rev. **C63**, 044002 (2001); A. Motzke et al., Phys. Rev. **C66**, 054002 (2002).